

A medium baseline search for $\nu_\mu \rightarrow \nu_e$ oscillations at a ν beam from muon decays

A. Bueno, M. Campanelli, A. Rubbia*
Institut für Teilchenphysik
ETHZ, CH-8093 Zürich, Switzerland

February 7, 2008

Abstract

The accurate knowledge of the $\bar{\nu}_e$ (ν_μ) beam produced in μ^- decays and the absence of ν_e ($\bar{\nu}_\mu$) contamination, make a future muon storage ring the ideal place to look for $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) oscillations. Using a detector capable of electron and muon identification with charge discrimination (e.g., the presently running NOMAD experiment), good sensitivities to $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) oscillations could be achieved. With the CERN-PS as a proton driver for a muon storage ring of the kind envisaged for a μ -collider, the LSND claim would be confirmed or disproved in a few years of running.

1 Introduction

In a recent study[1], we have investigated the possibility of performing a long baseline experiment using a neutrino beam from muon decays. In this paper, we concentrate on the possibility of a medium baseline experiment, motivated by the outstanding problem of the verification of the Los Alamos LSND experiment, which claims evidence for neutrino oscillations in the channels $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ [2] and $\nu_\mu \rightarrow \nu_e$ [3].

The latest KARMEN results[4] have come very close to contradicting the LSND claim, however the experimental sensitivity is marginal to conclusively exclude or confirm completely the LSND allowed solution.

The probability for neutrino oscillations between two families is given by

$$P(E_\nu) = \sin^2 2\theta \sin^2(1.27 \Delta m^2 \frac{L}{E_\nu}) \quad (1)$$

where θ is the mixing angle between the two neutrino flavours, and Δm^2 (in eV^2) is the difference of the squares of the neutrino masses.

The part of the LSND experiment solution with $\Delta m^2 > \approx 10 eV^2$ has recently been shown to be inconsistent with the data taken with the NOMAD detector[5] at the present CERN-WANF neutrino beam. In the NOMAD detector where electron identification capability is rather good and well understood, the current search for $\nu_\mu \rightarrow \nu_e$ oscillations[6] is limited in sensitivity by the systematic uncertainty of the prediction of the intrinsic ν_e component of the beam originating from primary K^+ , K_L^0 and μ^+ decays.

New experiments have been proposed to cover the $\Delta m^2 - \sin^2 2\theta$ regions indicated by the LSND claim:

*On leave from CERN, Geneva, Switzerland.

- ICARUS at Jura[7] would use the currently existing CERN WANF neutrino beam, extending the baseline to 17 km (allowing also tau appearance).
- At Fermilab, the MiniBOONE[8] experiment would address the low Δm^2 region by using a neutrino beam from the FNAL booster. The average energy is 1.5 GeV and the distance from the source to the detector is 1 km.
- The CERN/I216 letter of intent[9] suggested to reuse the old CERN-PS neutrino beam in order to obtain a low energy beam.

So far, MiniBOONE is the only approved experiment. To improve the oscillation result at low Δm^2 , the proponents of MiniBOONE have chosen an optimized L/E_ν in order to maximize the oscillation signal compared to backgrounds. This experiment should provide a strong signal in case the LSND effect is really due to neutrino oscillations. It suffers however from irreducible backgrounds associated to the intrinsic ν_e component of the beam and from misidentified electrons[8], that could make more problematic an interpretation of the observed effects.

We suggest that a complementary approach is to try to reduce background sources as much as possible, in order to obtain good statistical sensitivities, even in the case of small oscillation probabilities. We take advantage from the fact that when a neutrino beam is produced from muons of a definite sign, for example from the decay of negative muons¹, only one kind of flavor-antiflavor is produced, i.e.

$$\mu^- \rightarrow e^- \bar{\nu}_e \nu_\mu. \quad (2)$$

- In case of $\nu_\mu \rightarrow \nu_e$ oscillations, there will be appearance of ν_e neutrinos while the unoscillated beam contained only $\bar{\nu}_e$'s. Charge discrimination in the detector will trivially separate the two types of neutrino components:

$$\bar{\nu}_e + N \rightarrow e^+ + X \quad \text{intrinsic beam component,} \quad (3)$$

$$\nu_e + N \rightarrow e^- + X \quad \text{oscillated} \quad (4)$$

- In case of $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$ oscillations, there will be appearance of $\bar{\nu}_\mu$ neutrinos while the unoscillated beam contained only ν_μ 's.

$$\nu_\mu + N \rightarrow \mu^- + X \quad \text{intrinsic beam component,} \quad (5)$$

$$\bar{\nu}_\mu + N \rightarrow \mu^+ + X \quad \text{oscillated} \quad (6)$$

We stress that this experiment would test the transition $\nu_\mu \rightarrow \nu_e$ and its CP-conjugate $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ at the same time.

2 Event rates

We have already discussed our assumptions concerning the muon source in our previous paper[1] and recall here only the important points. Based on muon collider studies[10], we assume that the neutrino beam will be produced by the decay of a large quantity of stored muons[11]. The muons are produced in the decay chain of pions behind an appropriate target and are subsequently captured, cooled, accelerated and stored into a ring where they are let to decay. We compute the total number of muons accelerated in the following way:

$$N_\mu = f_{bunch} \times N_{p/bunch} \times Y_{\pi/p} \times Y_{\mu/\pi} \times t$$

where f_{bunch} is the bunch repetition rate, $N_{p/bunch}$ is the number of protons per bunch, $Y_{\pi/p}$ is the yield of pions per proton, and $Y_{\mu/\pi}$ is the yield of muons per pion.

¹In this paper we always refer to the decay of negative muons. The same considerations are obviously valid in the case of positive muons.

We need a proton driver in order to produce high energy protons for the muon source. The currently existing CERN-PS machine is able to accelerate² $f_{\text{bunch}} \times N_{p/\text{bunch}} \times t \approx 2.5 \times 10^{20}$ protons in a year at an energy of 19.2 GeV.

We assume that $Y_{\pi/p} \times Y_{\mu/\pi} = 0.08$ muon per proton will be trapped and accelerated into the storage ring[10]. With this assumption, the total number of muons accelerated is $2 \times 10^{19} \mu/\text{year}$. We consider that after 2 years of running, a total of $4 \times 10^{19} \mu$ will have decayed inside the storage ring. With an optimized geometrical configuration of the storage ring like a "cigare", in which two long straight sections are closed by two small arcs where the muons are strongly bent, half the muons will produce neutrinos in the right direction.

We chose a baseline from the CERN-PS to the CERN-Preveessin laboratory, close to the North Area. The distance from the PS is about 3.5 km (see Figure 1). We tentatively use the existing NOMAD detector as a target (total mass of 2.6 tons). It is presently located in the West Area and we assume that it can be moved to the North Area.

The muon beam energy is set to 7 GeV in order to obtain a relatively low energy neutrino beam. We compute the event rates assuming an unpolarized muon beam. The event rates for an integrated intensity of $4 \times 10^{19} \mu$'s are shown in Table 1. The energy spectra of ν_μ and $\bar{\nu}_e$ charged current events are shown in Figure 2. For comparison, we give the expected number of oscillated events for full mixing and various relevant Δm^2 parameters.

<i>No oscillations</i>	
ν_μ CC events	11600 μ^-
$\bar{\nu}_e$ CC events	5070 e^+
<i>With oscillations</i> $\Delta m^2 = 1\text{eV}^2, \sin^2 2\theta = 1$	
ν_e CC events	6600 e^-
$\bar{\nu}_\mu$ CC events	3300 μ^+
<i>With oscillations</i> $\Delta m^2 = 0.4\text{eV}^2, \sin^2 2\theta = 1$	
ν_e CC events	1580 e^-
$\bar{\nu}_\mu$ CC events	910 μ^+
<i>With oscillations</i> $\Delta m^2 = 0.2\text{eV}^2, \sin^2 2\theta = 1$	
ν_e CC events	430 e^-
$\bar{\nu}_\mu$ CC events	260 μ^+

Table 1: Neutrino event rates assuming the NOMAD detector (2.6 tons) as a target. The muon beam has an energy $E_\mu = 7$ GeV and the integrated intensity is $4 \times 10^{19} \mu$'s. The baseline is $L = 3.5$ km.

3 Results with the NOMAD detector

We have studied the possibility to perform $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$ and $\nu_\mu \rightarrow \nu_e$ oscillation appearance searches. The event selection is based on the identification of leading final state electrons or muons, and looks for opposite charges than the one expected in the unoscillated beam components.

We carried out the study of signal efficiency and background estimations using a full simulation of the events and the reconstruction packages currently used in NOMAD. All selection cuts are applied on the reconstructed quantities.

For the electron final state, the following backgrounds are the dominant ones:

- Charge confusion: $\nu_e + N \rightarrow e^+ + X \rightarrow "e^-"+X$

²See the discussion in Ref. [9] and we also stress that this number depends on a shared/dedicated mode of operation. We assume that neither LEP, nor LHC will be running during this time.

$\nu_\mu \rightarrow \nu_e$					
	ν_e CC (%)	ν_μ CC	ν_μ NC	$\bar{\nu}_e$ CC	$\bar{\nu}_e$ NC
Initial	100	11600	3867	5070	2028
Candidate highest P_T	59	196	99	1922	35.2
$P_{ele} > 2.5\text{GeV}$	41	4	2.1	1503	0.4
Calorimetry	35	0	0.7	1360	0.4
Charge	35	0	< 0.2	< 0.2	< 0.1

Table 2: Summary of detection efficiencies and expected backgrounds for a $\nu_\mu \rightarrow \nu_e$ oscillation search.

$\bar{\nu}_e \rightarrow \bar{\nu}_\mu$					
	$\bar{\nu}_\mu$ CC (%)	ν_μ CC	ν_μ NC	$\bar{\nu}_e$ CC	$\bar{\nu}_e$ NC
Initial	100	11600	3867	5070	2028
Candidate highest P_T	65	3874	2.5	0.5	0.4
Charge	65	2	1.1	0	0.18
Isolation	50	< 0.1	< 0.2	0	< 0.1

Table 3: Summary of detection efficiencies and expected backgrounds for a $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$ oscillation search.

- Asymmetrically reconstructed conversions or Dalitz decays: $\nu_\ell + N \rightarrow \nu_\ell + X$ and $X \rightarrow e^-(e^+) + X'$
- Misidentified hadrons: $\nu_\ell + N \rightarrow \nu_\ell + X$ and $h^- \rightarrow "e^-"$.

The NOMAD detector is well suited to search for electron charged current events. The electron identification is good and well understood. Charge separation of leading electron is also very well known. Charge confusion arises primarily from early bremsstrahlung with leading photon converting asymmetrically into a e^+e^- pair where the e^- is lost. This process is rare and the simulation gives in the low energy region a contamination at the level of one in ten thousand.

An important source of background for the electron search comes from Dalitz decays and π^0 conversions (where the γ converts close to the primary interaction vertex). Those non-prompt electrons, embedded in the recoiling hadronic system, are in general less isolated than the prompt ones. Hence, an efficient rejection of this background is obtained by demanding that the electron candidate has the highest transverse momentum in the event. Additional discrimination is achieved by means of a lower cut on the measured electron momentum. Remnant backgrounds, due to hadron misidentification, can be reduced to a negligible level by comparing the reconstructed electron momentum (tracking) to the associated measured energy (calorimetry).

Table 2 summarizes the results obtained for electrons, when normalized to 4×10^{19} muons. For an overall 35% electron detection efficiency (ϵ_{e^-}), the expected background is well below one event.

For the muon search, we use the standard muon identification algorithm used in the NOMAD experiment. It has a momentum threshold of about 3 GeV imposed by the requirement of reachability of the muon chambers. After the cuts listed in Table 3, the detection efficiency for μ^+ (ϵ_{μ^+}) is 50 %. The background contamination for the $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$ oscillation search is negligible. The small punch-through background is completely suppressed by a kinematical isolation requirement, since a misidentified muon will come from the jet.

In case of positive signal, the simultaneous use of both channels provides a test of CP

conservation in the leptonic system. Figure 3 shows a candidate event for the $\nu_\mu \rightarrow \nu_e$ oscillation. The signal events appear as remarkably clean and simple to reconstruct.

In case of negative result, combining both channels assuming CP conservation, the 90%C.L. limit on $\nu_\mu \rightarrow \nu_e$ oscillations, for a total flux of 4×10^{19} muons, can be obtained from the following:

$$P(\nu_\mu \rightarrow \nu_e) < \frac{2.3}{11600 \times \epsilon_{e-} + 5070 \times \epsilon_{\mu+}} = 3.5 \times 10^{-4} \quad (7)$$

This corresponds to a limit on the mixing angle for large Δm^2 of

$$\sin^2 2\theta < 7 \times 10^{-4} \quad 90\%C.L. \quad (8)$$

and $\Delta m^2 > 2 \times 10^{-2} \text{ eV}^2$ for maximal mixing. The 90% C.L. exclusion contour is shown in Figure 4 compared to the LSND positive solution, the current limits from NOMAD and CCFR, and the expected limit from MiniBOONE.

4 Conclusions

Testing the muon yield per proton sent on a target, is the first step in the R&D program towards the construction of a future muon collider. At CERN, the CERN-PS would be the natural proton source for this device. We have shown that with this setup, and with the possibility of recycling parts of an already existing detector (NOMAD), it would be possible to perform an experiment to increase the present limits on $\nu_\mu - \nu_e$ oscillations. The area in the parameter space indicated by the LSND result can be fully covered, with a sensitivity compared to that of the MiniBOONE experiment, but with a completely different approach.

References

- [1] A. Bueno, M. Campanelli and A. Rubbia, hep-ph/9808485, August 1998.
- [2] C. Athanassopoulos *et al.*, (LSND Collaboration), *Phys. Rev. C* **54**, 2685 (1996); C. Athanassopoulos *et al.*, (LSND Collaboration), *Phys. Rev. Lett.* **77**, 3082 (1996); *Phys. Rev. Lett.* **75**, 2650 (1995). Updated results including 1997 data can be found at <http://www.neutrino.lanl.gov/LSND/papers.html>.
- [3] C. Athanassopoulos *et al.*, (LSND Collaboration), LA-UR-97-1998/UCRHEP-E191, submitted to *Phys. Rev. C*; C. Athanassopoulos *et al.*, *Phys. Rev. Lett.* **81**, 1774 (1998).
- [4] B. Zeitnitz, KARMEN Collab., proceedings of the Neutrino98 Conference, Takayama, Japan, June 1998.
- [5] NOMAD Collab, J. Altegoer *et al.*, *Nucl. Instr. and Meth. A* **404** (1998) 96.
A. Rubbia, *Nucl. Phys. B(Proc. Suppl.)* **40** (1995) 93.
- [6] V. Valuev, NOMAD Collaboration, proceedings of the International Conference on High Energy Physics, Jerusalem, Israel, 19-26 August 1997, to be published.
V. Valuev, “*Recherches des oscillations de neutrinos $\nu_\mu \rightarrow \nu_e$ dans l’expérience NOMAD*”; Ph.D. Thesis Université de Paris VII, Denis-Diderot, 1998.
A. Bueno *et al.*, NOMAD internal memo # 98-016.
- [7] ICARUS-CERN-Milano Coll., CERN/SPSLC 96-58, SPSLC/P 304, December 1996;
J. P. Revol *et al.*, ICARUS-TM-97/01, 5 March 1997, unpublished.
- [8] Updated information on the BOONE proposal can be found at <http://www.neutrino.lanl.gov/BooNE/>.
- [9] N. Armenise *et al.*, CERN-SPSC/97-21, October 1997.
- [10] C.M. Ankenbrandt *et al.*, Muon Collider Collab., “Status Report”, see <http://www.cap.bnl.gov/mumu/>
- [11] S. Geer, *Phys. Rev. D* **57**, 6989 (1998).




Figure 1: Schematic overview of the CERN sites.




Figure 2: Predicted muon neutrino and electron anti-neutrino charged current event energy distribution for an integrated statistics of 4×10^{19} muons.




Figure 3: A ν_e CC oscillated signal event in NOMAD. The leading particle is 5 GeV electron stopping in the Electromagnetic Calorimeter




Figure 4: Predicted 90%C.L. limit for this experiment compared to expected MiniBOONE limit for neutrino oscillation appearance. Negative results from previous NOMAD, CCFR and reactor experiments are also shown.